

Semileptonic B Decays with $N_f = 2 + 1$ Dynamical Quarks

J.Shigemitsu^a, C.T.H.Davies^b, A.Dougall^b, K.Foley^c, E.Gamiz^b, A.Gray^a, E.Gulez^a, G.P.Lepage^c, M.Wingate^d

^aPhysics Department, The Ohio State University, Columbus, OH 43210, USA.

^bDepartment of Physics & Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK.

^cLaboratory of Elementary Particle Physics, Cornell University, Ithaca, NY 14853, USA.

^dInstitute for Nuclear Theory, University of Washington, Seattle, WA 98115, USA.

Semileptonic, $B \rightarrow \pi l \bar{\nu}$, decays are studied on the MILC dynamical configurations using NRQCD heavy and Asqtad light quarks. We work with light valence quark masses ranging between m_s and $m_s/8$. Preliminary simple linear chiral extrapolations have been carried out for form factors $f_{\parallel}$ and $f_{\perp}$ at fixed E_{π} . The chirally extrapolated results for the form factors $f_+(q^2)$ and $f_0(q^2)$ are then fit to the Becirevic-Kaidalov (BK) ansatz. Preliminary estimates of the CKM matrix element $|V_{ub}|$ are presented based on recently published branching fractions for $B^0 \rightarrow \pi^- l^+ \nu$ exclusive decays by the CLEO collaboration.

1. Introduction

First principles calculations of B meson semileptonic decay form factors are crucial for determining the CKM matrix elements $|V_{cb}|$ and $|V_{ub}|$. Recent progress on the lattice towards this goal comes from two major developments: the ability to go beyond the quenched approximation with close to realistic dynamical quark content [1,2] and the use of improved staggered light quarks in heavy-light simulations [3]. We report here on unquenched studies of $B \rightarrow \pi, l \bar{\nu}$ decays on the lattice using one of the coarse MILC $N_f = 2 + 1$ dynamical sets [1], NRQCD b quarks and improved staggered (Asqtad) light quarks. The light dynamical quark mass is fixed at $m_{dyn} = m_s/4$ and we vary the light valence quark mass between m_s and $m_s/8$.

2. Form Factors

Semileptonic form factors parameterise the hadronic matrix elements of electroweak currents between a B meson and a π or a ρ . In particular,

one has

$$\langle \pi | V^{\mu} | B \rangle = f_+(q^2) \left[p_B^{\mu} + p_{\pi}^{\mu} - \frac{M_B^2 - m_{\pi}^2}{q^2} q^{\mu} \right] + f_0(q^2) \frac{M_B^2 - m_{\pi}^2}{q^2} q^{\mu} \quad (1)$$

$$= \sqrt{2M_B} [v^{\mu} f_{\parallel} + p_{\perp}^{\mu} f_{\perp}] \quad (2)$$

with

$$v^{\mu} = \frac{p_B^{\mu}}{M_B}, \quad p_{\perp}^{\mu} = p_{\pi}^{\mu} - (p_{\pi} \cdot v) v^{\mu}, \quad q^{\mu} = p_B^{\mu} - p_{\pi}^{\mu}.$$

A lattice calculation of the relevant matrix element starts with the three-point correlator

$$C^{(3)}(\vec{p}_{\pi}, \vec{p}_B, t, T_B) = \sum_{\vec{z}} \sum_{\vec{y}} \langle \Phi_{\pi}(0) V_{lat}^{\mu}(\vec{z}, t) \Phi_B^{\dagger}(\vec{y}, T_B) \rangle \times e^{i\vec{p}_B \cdot \vec{y}} e^{i(\vec{p}_{\pi} - \vec{p}_B) \cdot \vec{z}} \quad (3)$$

where Φ_{π} and Φ_B are interpolating operators for the π and B mesons respectively. The three-point correlator is fit to the form

$$C^{(3)}(\vec{p}_{\pi}, \vec{p}_B, t, T_B) \rightarrow \sum_{k=0}^{N_{\pi}-1} \sum_{j=0}^{N_B-1} (-1)^{k*(t-1)} (-1)^{j*(T_B-t)} \times A_{j,k} e^{-E_{\pi}^{(k)}(t-1)} e^{-E_B^{(j)}(T_B-t)} \quad (4)$$




Figure 1. The form factors $f_+(q^2)$ and $f_0(q^2)$, extrapolated to the physical pion. Only statistical errors are shown. The full curves correspond to a Becirevic-Kaidalov (BK) parametrization fit to the data (see text).

and the goal is to extract the ground state contribution A_{00} . Working in the B meson rest frame, the ground state amplitudes $A_{00}(V^\mu)$ are related to the form factors $f_\parallel$ and $f_\perp$ in a simple way.

$$f_\parallel = \frac{A_{00}(V^0)}{\sqrt{\zeta_\pi^{(0)} \zeta_B^{(0)}}} \sqrt{2E_\pi} Z_{V_0} \quad (5)$$

$$f_\perp = \frac{A_{00}(V^k)}{\sqrt{\zeta_\pi^{(0)} \zeta_B^{(0)}}} \sqrt{2E_\pi} Z_{V_k} p_\pi^k \quad (6)$$

where $\zeta_{\pi,B}^{(0)}$ are the groundstate amplitudes of the π and B two-point correlators respectively and Z_{V_μ} are the matching factors between the lattice heavy-light current V_{lat}^μ and the current in the continuum theory. We use one-loop estimates for these matching factors [4].

3. The Form Factors $f_+(q^2)$ and $f_0(q^2)$ at the Physical Pion

Although our simulations have been carried out with light quark masses as low as $m_s/8$ one still needs to extrapolate the form factors, determined above, to the physical pion. To date, we have only carried out simple linear chiral extrapolations. We first interpolate $f_\parallel$ and $f_\perp$ to common values of E_π , the pion energy in the B rest frame. These are then extrapolated linearly to the physical pion for several fixed values of E_π .

From the chirally extrapolated $f_\parallel$ and $f_\perp$ one obtains the form factors $f_+(q^2)$ and $f_0(q^2)$ at the physical pion. This is shown as the circles in Fig.1. Our results are currently limited to the $q^2 \geq 15 \text{ GeV}^2$ region. Recently, a very promising approach to low q^2 form factors has been developed, namely “Moving NRQCD”, which will allow us to overcome this limitation [5,6]. In the mean time, however, we will rely on a model ansatz to extend our form factor results into the low q^2 regime. Specifically, we employ an ansatz introduced by Becirevic & Kaidalov (BK) [7],

$$f_+(q^2) = \frac{C_B (1 - \alpha_B)}{(1 - \tilde{q}^2)(1 - \alpha_B \tilde{q}^2)} \quad (7)$$

$$f_0(q^2) = \frac{C_B (1 - \alpha_B)}{(1 - \tilde{q}^2/\beta_B)} \quad (8)$$

($\tilde{q}^2 \equiv q^2/M_{B^*}^2$). This ansatz satisfies the kinematic constraint $f_+(0) = f_0(0)$, HQET scaling laws and the requirement of a pole in $f_+(q^2)$ at $q^2 = M_{B^*}^2$. We find an excellent fit to this BK ansatz using the physical M_{B^*} mass and this is shown as the full curves in Fig.1 (a satisfactory BK parametrization was not possible before the chiral extrapolation). The fit parameters are

$$C_B = 0.42(3), \quad \alpha_B = 0.41(7), \quad \beta_B = 1.18(5),$$

which translates into

$$f_0(0) = f_+(0) = 0.251(15)$$

and an effective pole in $f_0(q^2)$ at $q^2 = (M_{f_0}^{pole})^2 = 33.35(1.36) \text{ GeV}^2$. Both $f_{0,+}(0)$ and $M_{f_0}^{pole}$ are in good agreement with a recent semileptonic B decay analysis based on Sum Rules [8]. The data

points in Fig.1 and the $f_{0,+}(0)$ quoted above include only statistical and fitting errors. Further systematic errors are discussed in the next section. A comparison of our new dynamical form factor results with old quenched data is given, for instance, in reference [9].

4. Estimating $|V_{ub}|$

The CLEO collaboration has published branching fractions for exclusive semileptonic B decays, including binning into several q^2 ranges [10]. We combine these experimental inputs with lattice results for $f_+(q^2)$ to extract values for the CKM matrix element $|V_{ub}|$. The differential decay rate for $B^0 \rightarrow \pi^-, l^+ \nu$,

$$\frac{1}{|V_{ub}|^2} \frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} p_\pi^3 |f_+(q^2)|^2 \quad (9)$$

can be integrated to give $\frac{\Gamma}{|V_{ub}|^2}$ and the partial width Γ can be determined from CLEO's branching fraction and the Particle Data Group's B^0 lifetime of $1.542 \pm 0.016 ps$. Our preliminary estimate for $|V_{ub}|$ is then

$$|V_{ub}| = \begin{cases} 3.86(32)(58) \times 10^{-3} & 0 \leq q^2 \leq q_{max}^2 \\ 3.52(73)(44) \times 10^{-3} & 16 GeV^2 \leq q^2 \end{cases}$$

where the two values correspond to either using the entire allowed q^2 range or restricting both experiment and theory to the $q^2 \geq 16 GeV^2$ region. The first error is experimental and the second is our current best estimate of lattice statistical and systematic errors added in quadrature. In addition to $4 \sim 6\%$ statistical errors, we estimate $\sim 9\%$ higher order perturbative matching, $\sim 5\%$ chiral extrapolation, $\sim 5\%$ relativistic and discretization errors. This adds up to $\sim 12.5\%$ lattice errors for $|V_{ub}|$ obtained from the $q^2 \geq 16 GeV^2$ region. For $|V_{ub}|$ based on the full q^2 range we increase the lattice errors to 15% (an additional 8% added in quadrature) taking into account the need to rely on the BK parametrization to enter the low q^2 region.

5. Summary

Unquenched simulations of heavy meson semileptonic decays are now feasible and we re-

port here on the first such calculations using NRQCD heavy and Asqtad light quarks (see talk by Okamoto for results using Fermilab heavy quarks [11]). The use of the improved staggered light quark action has allowed for significantly smaller statistical and chiral extrapolation errors than in the past. Combining lattice results for $f_+(q^2)$ with experimental branching fraction data has led to preliminary estimates of $|V_{ub}|$.

Many improvements are planned: inclusion of all dimension 4 ($1/M$, α_s/M and $a\alpha_s$) current corrections, more sophisticated chiral extrapolations [12], and simulations at other dynamical quark masses and lattice spacings. Use of "Moving NRQCD" [5] will also allow us to simulate directly at lower q^2 .

Acknowledgements : This work was supported by the DOE, PPARC and NSF. Simulations were carried out at NERSC.

REFERENCES

1. C.W.Bernard *et al.* (MILC); Phys.Rev. **D64**, 054506 (2001).
2. C.T.H.Davies *et al.* (MILC,HPQCD,UKQCD); Phys.Rev.Lett. **92**, 022001 (2004).
3. M. Wingate *et al.*; Phys. Rev. **D67**, 054505 (2003).
4. E.Gulez; these proceedings; E.Gulez *et al.*; Phys.Rev. **D69**, 074501 (2004).
5. K.Foley; these proceedings
K.Foley and G.P.Lepage; Nucl.Phys. Proc. Suppl.**119**, 635 (2003).
6. J.Sloan; Nucl.Phys. Proc. Suppl.**63**, 365 (1998)
S.Hashimoto and H.Matsufuru; Phys.Rev. **D54**, 4578 (1996).
7. D.Becirevic and A.B.Kaidalov; Phys. Lett. **B478**, 417 (2000).
8. P.Ball and R.Zwicky; hep-ph/0406232.
9. A.Gray; hep-ph/0407229.
10. S.B.Athar *et al.* (CLEO); Phys.Rev. **D68**, 072003 (2003), [hep-ex/0304019].
11. M.Okamoto; these proceedings
12. C.Aubin and C.Bernard; these proceedings and private communication.